

graviton

$$J = 2$$

graviton MASS

It is likely that the graviton is massless. More than fifty years ago Van Dam and Veltman (VANDAM 70), Iwasaki (IWASAKI 70), and Zakharov (ZAKHAROV 70) almost simultaneously showed that in the linear approximation a theory with a finite graviton mass does not approach GR as the mass approaches zero. Attempts have been made to evade this "vDVZ discontinuity" by invoking modified gravity or nonlinear theory by De Rahm (DE-RHAM 17) and others. More recently, the analysis of gravitational wave dispersion has led to bounds that are largely independent of the underlying model, even if not the strongest. We quote the best of these as our best limit.

Experimental limits have been set based on a Yukawa potential (YUKA), dispersion relation (DISP), or other modified gravity theories (MGRV).

The following conversions are useful: $1 \text{ eV} = 1.783 \times 10^{-33} \text{ g} = 1.957 \times 10^{-6} m_e$; $\lambda_C = (1.973 \times 10^{-7} \text{ m}) \times (1 \text{ eV}/m_g)$.

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$<1.76 \times 10^{-23}$	1 ABBOTT	21	DISP LIGO Virgo catalog GWTC-2
$<3.2 \times 10^{-23}$	2 BERNUS	20	YUKA Planetary ephemeris INPOP19a
$<2 \times 10^{-28}$	3 SHAO	20	DISP Binary pulsar Galileon radiation
$<4.7 \times 10^{-23}$	4 ABBOTT	19	DISP LIGO Virgo catalog GWTC-1
$<7 \times 10^{-23}$	5 BERNUS	19	YUKA Planetary ephemeris INPOP17b
$<3.1 \times 10^{-20}$	6 MIAO	19	DISP Binary pulsar orbital decay rate
$<1.4 \times 10^{-29}$	7 DESAI	18	YUKA Gal cluster Abell 1689
$<5 \times 10^{-30}$	8 GUPTA	18	YUKA Using SPT-SZ
$<3 \times 10^{-30}$	8 GUPTA	18	YUKA Using Planck all-sky SZ
$<1.3 \times 10^{-29}$	8 GUPTA	18	YUKA Using redMaPPer SDSS-DR8
$<6 \times 10^{-30}$	9 RANA	18	YUKA Weak lensing in massive clusters
$<8 \times 10^{-30}$	10 RANA	18	YUKA SZ effect in massive clusters
$<1.0 \times 10^{-23}$	11 WILL	18	YUKA Perihelion advances of planets
$<7 \times 10^{-23}$	4 ABBOTT	17	DISP Combined dispersion limit from three BH mergers
$<1.2 \times 10^{-22}$	4 ABBOTT	16	DISP Combined dispersion limit from two BH mergers
$<2.9 \times 10^{-21}$	12 ZAKHAROV	16	YUKA S2 star orbit
$<5 \times 10^{-23}$	13 BRITO	13	MGRV Spinning black holes bounds
$<6 \times 10^{-32}$	14 GRUZINOV	05	MGRV Solar System observations
$<6 \times 10^{-32}$	15 CHOUDHURY	04	YUKA Weak gravitational lensing
$<9.0 \times 10^{-34}$	16 GERSHTEIN	04	MGRV From Ω_{tot} value assuming RTG
$<8 \times 10^{-20}$	17,18 FINN	02	DISP Binary pulsar orbital period decrease
$<7 \times 10^{-23}$	TALMADGE	88	YUKA Solar system planetary astrometric data
$<1.3 \times 10^{-29}$	19 GOLDHABER	74	YUKA Rich clusters
$<7 \times 10^{-28}$	HARE	73	YUKA Galaxy
$<8 \times 10^4$	HARE	73	YUKA 2γ decay

- ¹ ABBOTT 21 assumed modified gravitational-wave dispersion to establish a limit on graviton mass, using LIGO-Virgo O1-O3a binary black hole (BBH) events.
- ² BERNUS 20 use the latest solution of the ephemeris INPOP (19a) in order to improve the constraint in BERNUS 19 on the existence of a Yukawa suppression to the Newtonian potential, generically associated to a gravitons mass.
- ³ SHAO 20 sets limit, 95% CL, based on non-observation of excess gravitational radiation in 14 well-timed binary pulsars in the context of the cubic Galileon model.
- ⁴ ABBOTT 19, ABBOTT 17, and ABBOTT 16 assumed modified gravitational waves dispersion to establish limits on graviton mass.
- ⁵ BERNUS 19 use the planetary ephemeris INPOP 17b to constraint the existence of a Yukawa suppression to the Newtonian potential, generically associated to a gravitons mass.
- ⁶ MIAO 19 90% CL limit is based on orbital period decay rates of 9 binary pulsars using a Bayesian prior uniform in graviton mass. Limit becomes $< 5.2 \times 10^{-21}$ eV for a prior uniform in $\ln(m_g)$.
- ⁷ DESAI 18 limit based on dynamical mass models of galaxy cluster Abell 1689.
- ⁸ GUPTA 18 obtains graviton mass limits using stacked clusters from 3 disparate surveys.
- ⁹ RANA 18 limit, 68% CL, obtained using weak lensing mass profiles out to the radius at which the cluster density falls to 200 times the critical density of the Universe. Limit is based on the fractional change between Newtonian and Yukawa accelerations for the 50 most massive galaxy clusters in the Local Cluster Substructure Survey. Limits for other CL's and other density cuts are also given.
- ¹⁰ RANA 18 limit, 68% CL, obtained using mass measurements via the SZ effect out to the radius at which the cluster density falls to 500 times the critical density of the Universe for 182 optically confirmed galaxy clusters in an Atacama Cosmology Telescope survey. Limits for other CL's and other density cuts are also given.
- ¹¹ WILL 18 limit from perihelion advances of the planets, notably Earth, Mars, and Saturn. Alternate analysis yields $< 6 \times 10^{-24}$.
- ¹² ZAKHAROV 16 constrains range of Yukawa gravity interaction from S2 star orbit about black hole at Galactic center. The limit is $< 2.9 \times 10^{-21}$ eV for $\delta = 100$.
- ¹³ BRITO 13 explore massive graviton (spin-2) fluctuations around rotating black holes.
- ¹⁴ GRUZINOV 05 uses the DGP model (DVALI 00) showing that non-perturbative effects restore continuity with Einstein's equations as the graviton mass approaches zero, then bases his limit on Solar System observations.
- ¹⁵ CHOUDHURY 04 concludes from a study of weak-lensing data that masses heavier than about the inverse of 100 Mpc seem to be ruled out if the gravitation field has the Yukawa form.
- ¹⁶ GERSHTEIN 04 use non-Einstein field relativistic theory of gravity (RTG), with a massive graviton, to obtain the 95% CL mass limit implied by the value of $\Omega_{tot} = 1.02 \pm 0.02$ current at the time of publication.
- ¹⁷ FINN 02 analyze the orbital decay rates of PSR B1913+16 and PSR B1534+12 with a possible graviton mass as a parameter. The combined frequentist mass limit is at 90%CL.
- ¹⁸ As of 2020, limits on dP/dt are now about 0.1% (see T. Damour, "Experimental tests of gravitational theory," in this *Review*).
- ¹⁹ GOLDBABER 74 establish this limit considering the binding of galactic clusters, corrected to Planck $h_0 = 0.67$.

graviton REFERENCES

ABBOTT	21	PR D103 122002	R. Abbott <i>et al.</i>	(LIGO and Virgo Collabs.)
BERNUS	20	PR D102 021501	L. Bernus <i>et al.</i>	
SHAO	20	PR D102 024069	L. Shao, N. Wex, S.-Y. Zhou	
ABBOTT	19	PR D100 104036	B.P. Abbott <i>et al.</i>	(LIGO and Virgo Collabs.)
BERNUS	19	PRL 123 161103	L. Bernus <i>et al.</i>	
MIAO	19	PR D99 123015	X. Miao, L. Shao, B.-Q. Ma	
DESAI	18	PL B778 325	S. Desai	(HYDER)
GUPTA	18	ANP 399 85	S. Gupta, S. Desai	

RANA	18	PL B781 220	A. Rana <i>et al.</i>	(DELHI)
WILL	18	CQG 35 17LT01	C.M. Will	
ABBOTT	17	PRL 118 221101	B.P. Abbott <i>et al.</i>	(LIGO and Virgo Collabs.)
DE-RHAM	17	RMP 89 025004	C. de Rham <i>et al.</i>	
ABBOTT	16	PRL 116 061102	B.P. Abbott <i>et al.</i>	(LIGO and Virgo Collabs.)
ZAKHAROV	16	JCAP 1605 045	A.F. Zakharov <i>et al.</i>	
BRITO	13	PR D88 023514	R. Brito, V. Cardoso, P. Pani	(LISB, MISS, HSCA+)
GRUZINOV	05	NAST 10 311	A. Gruzinov	(NYU)
CHOUDHURY	04	ASP 21 559	S.R. Choudhury <i>et al.</i>	(DELPH, MELB)
GRSHTSTEIN	04	PAN 67 1596	S.S. Gershtein <i>et al.</i>	(SERP)
		Translated from YAF 67 1618.		
FINN	02	PR D65 044022	L.S. Finn, P.J. Sutton	
DVALI	00	PL B485 208	G.R. Dvali, G. Gabadadze, M. Porrati	(NYU)
TALMADGE	88	PRL 61 1159	C. Talmadge <i>et al.</i>	(JPL)
GOLDHABER	74	PR D9 1119	A.S. Goldhaber, M.M. Nieto	(LANL, STON)
HARE	73	CJP 51 431	M.G. Hare	(SASK)
IWASAKI	70	PR D2 2255	Y. Iwasaki	
VANDAM	70	NP B22 397	H. van Dam, M. Veltman	(UTRE)
ZAKHAROV	70	JETPL 12 312	V.I. Zakharov <i>et al.</i>	
